

Evaluation of Serum Level of Endoglin in Psoriatic Patients

M.A.Ibrahim¹, E.M.Akl¹, N.A.Abdul-Hafeez² and S.Z.El-Shebl¹

¹Dermatology, Venereology Dept., Faculty of Medicine, Benha Univ., Benha, Egypt

²clinical pathology and immunology, Dept., Faculty of Medicine, Benha Univ., Benha, Egypt

E.Mail: sarahelshebl@gmail.com

Abstract

Psoriasis is a lifelong chronic inflammatory disease of the skin with an estimated prevalence of 1-3% in the community. Endoglin (Eng; CD105) is an auxiliary receptor for the transforming growth factor β (TGF- β) family of proteins and has been shown to be essential for angiogenesis and is predominantly expressed on proliferating endothelial cells. Its mutation leads to hereditary hemorrhagic telangiectasia. The aim of this study is to evaluate the serum level of endoglin in patients with psoriasis and its correlation with the disease severity. Endoglin level might have been assessed in the serum from claiming 54 patients and 30 control subjects utilizing elisa. Information examination might have been conveyed crazy for the whole companion. Facts were performed utilizing programming. Comes about from claiming this ponder demonstrated that those intend serum level from claiming endoglin might have been higher On psoriatic patients over control one assembly for a huge quality. Our discoveries recommend that, there may be a part of endoglin in the pathophysiology Furthermore seriousness of psoriasis.

Key words: Psoriasis, Endoglin.

1. Introduction

Psoriasis may be a standout amongst the The greater part regular mankind's skin ailments Also may be recognized with bring magic hereditary underpinnings. It will be described by over the top Growth and aberrant separation about keratinocytes, Anyhow may be completely reversible for suitable treatment [1].

Psoriasis is portrayed Eventually Tom's perusing perceptible (sharply demarcated, red What's more somewhat brought up lesions with silver whitish scales) and comparing minute (acanthosis, papillomatosis, parakeratosis What's more munro micro abscess in the stratum corneum) skin alterations and prompts significant hindrance of the personal satisfaction of term of the influenced patients [2].

Clinical phenotypes from claiming psoriasis incorporate psoriasis vulgaris, guttate psoriasis, opposite psoriasis, erythrodermic psoriasis and pustular psoriasis. Separate clinical phenotypes of the ailment might a chance to be watched Significantly in the same tolerant in the course of the ailment progression [3].

The pathogenesis of psoriasis includes dynamic cooperations the middle of numerous Mobile sorts Furthermore various cytokines because of the opposition on triggers, culminating in the disturbance for skin resistant homeostasis clinched alongside hereditarily predisposed people [4].

Medicines for psoriasis incorporate topicals for example, such that corticosteroids, and also operators for example, anthralin, manufactured Vitamin D3, What's more Vitamin A, phototherapy including wide Furthermore narrowband-UVB, laser UVB, and psoralen UVA (PUVA); systemics for example, cyclosporine, methotrexate, and retinoid receptor inhibitors (Acitretin), and living therapeutics including TNF- α inhibitors and cytokine inhibitors for example, anti-IL23p40 Also IL-17 inhibitors [5].

Traditionally, psoriasis need been viewed Likewise An malady influencing main the skin Also joints. To later years, investigations from distinctive nations have demonstrated that psoriasis will be a systemic incendiary disease, which will be frequently connected for Different

comorbidities. To particular, there is a more excellent danger for Creating extreme vascular occasions for example, cardiovascular Furthermore cerebrovascular maladies [6].

Those pervasiveness rates about cardiovascular danger figures need aid expanded over psoriasis patients, including hypertension, diabetes, obesity, dyslipidemia, subclinical atherosclerosis, Furthermore smoking auto. It need been recommended that systemic aggravation might give acceptable An robotic join the middle of psoriasis Furthermore cardiometabolic issue [7].

Patients with psoriasis have a higher occurrence for metabolic syndrome over all population, Furthermore patients for a a greater amount extreme manifestation of psoriasis need a higher occurrence about metabolic syndrome over the individuals for An gentle type [8].

Those association between psoriasis and corpulence will be most likely bidirectional, for stoutness predisposing will psoriasis and psoriasis favouring stoutness [9].

Various instruments might make included for demonstrating those connection the middle of psoriasis and stoutness. Patients for psoriasis might bring a higher hazard for social isolation, poor consuming habits, depression, diminished physical activity, What's more expanded liquor utilization [10].

Endoglin (Eng; CD105) is a transmembrane glycoprotein, which is communicated Eventually Tom's perusing endothelial units (EC), Anyhow not Toward platelets, and assumes a basic physiological part in the cardiovascular framework [11]. Different locales from claiming outflow incorporate syncytiotrophoblasts from claiming full-term placenta, stromal units Furthermore certain haemopoietic phones [12]. It may be an assistant receptor for those transforming Growth figure β (TGF- β) gang for proteins Also need been indicated to be fundamental for angiogenesis [13].

Human endoglin will be a180 kDa homo-dimeric protein of 658 amino residues that holds a extracellular domain, An single transmembrane space and An short intracellular space [14].

Endoglin may be structurally identified with

betaglycan, another TGF- β kind iii assistant receptor (T β RIII), for helter skelter similitude in the transmembrane and intracellular districts [15].

The human endoglin gene will be found around human chromosome 9 with area of the cytogenic band toward 9q34. 11. Endoglin glycoprotein is encoded Eventually Tom's perusing 39,757 bp Furthermore makes as 658 aminic acids [16].

It will be communicated on human pre-erythroblasts, macrophages, leukemic. Units of the lymphoid Furthermore myeloid lineages, Furthermore In higher levels with respect to syncytiotrophoblasts about expression placenta Furthermore vascular endothelial units [17].

Endoglin will be communicated In low levels for resting endothelial cells, Yet it is profoundly communicated to vascular endothelial units at locales of animated angiogenesis for example, tumor vessels, aggravated tissues, recuperating wounds, psoriatic skin, aggravated synovial arthritis, upon vascular injury, What's more Throughout embryogenesis [18].

Transformations in mankind's endoglin looking into chromosome 9q34ter bring about the autosomal prevailing structure about inherited hemorrhagic telangiectasia sort 1, an uncommon state described Toward epistaxis Also vascular abnormality [19].

Supporting the connection the middle of Eng and metabolism, An association the middle of plasma levels for Eng Furthermore glycemia might have been discovered Previously, diabetic patients [20]. In addition, endoglin lack will be identified with endothelial brokenness What's more there will be an acceptable companionship between endothelial brokenness Also alterations done glucose digestion system alternately metabolic syndrome [21].

2. Materials and methods

2.1 The Study Population

This will be a case-control study. Those current ponder required been directed the middle of october 2019 What's more february 2019 after those Regard Toward examination council during employees of Medicine, Benha school. The ponder included 84 subjects that were haphazardly chose from the individuals going to the dermatology Outpatient's facility toward Benha school Hospital, staff about Medicine, Benha University, Qalyobia, egypt. Former will start of the study, each liable might have been educated over the point of the study What's more an educated assent might have been acquired starting with every singular preceding example accumulation. They were ordered under three aggregations; those to start with aggregation included 54 patients enduring from psoriasis; and the secod gathering included 30 Obviously solid subjects for matched agdistis and sex Concerning illustration control aggregation.

The sum contemplated subjects were tried for the serum level for endoglin.

Only the cases were tested for:

- 1- Lipid profile (HDL,LDL,cholesterol and triglecrides).
- 2- Fasting blood sugar.
- 3- Complete blood picture (CBC).
- 4- Serum albumin.
- 5- Erythrocyte sedimentation rate (ESR).
- 6- C-reactive protein (CRP).

2.2 ELISA assays of endoglin

Quantitative determinations of serum endoglin were achieved using a sandwich enzyme-linked immune-sorbent assay technology (ELISA) was used to detect serum level of endoglin using a commercial Human endoglin ELISA Kit for research use only (Cloud-Clone/Houston, Txs, USA).

2.3 Statistical analysis

Those gathered information might have been revised, coded, tabulated Furthermore acquainted utilizing the PC program SPSS (Statistical bundle for social science) adaptation 20. Information were exhibited Also suitability examination might have been carried as stated by the sort of information gotten for every parameter. Learner t test might have been used to analyze intend for two gatherings from claiming quantitative information. For the correlation from claiming intend for more than two Assemblies about quantitative information restricted examination of fluctuation (ANOVA) might have been utilized. Chi-Square test might have been used to analyze the relationship between two qualitative variables. Fisher's accurate test: might have been used to analyze the association the middle of two qualitative variables At the needed number is less 5 in more than 20% of phones. Connection analysis: with survey those quality from claiming cooperation the middle of two quantitative variables. The relationship coefficient characterizes those quality What's more bearing of the straight relationship between two variables. P may be huge Assuming that <0.05 during certainty interim 95%.

3. Results

3.1 The Study Population

The current study included 54 patients (35 males and 19 females). The average age of psoriasis patients was 52.02 ± 13.09 years. Male patients represented 64.8%, while females represented 35.2%. Mean disease duration was 15.35 ± 12.73 years. All patients had gradual onset, progressive course. The study included another ssecond group consisting of 30 healthy age and sex matched subjects (13 males and 17females) as a control group with an average age 47.67 ± 10.18 years .

Smoking habit found in 17(31.5%) and 8 (26.7%) in case and control group respectively.

By comparison between two groups, there was no significant difference regarding sex ($p=0.057$) ,age ($p=0.12$) , BMI ($p=0.85$) or smoking habit ($p=0.64$) Table (1).

Table (1) Personal characteristics of the studied groups.

	Case group (54)		Control group (30)		Statistical test	P value
	No	%	No	%		
Gender						
Male	35	64.8	13	43.3	X ² = 3.63	0.057
Female	19	35.2	17	56.7		
Age/years	52.02±13.09 (22-75)		47.67±10.18(31-66)		St t=1.57	0.12
Mean ±SD (range)						
BMI	28.59±3.62 (20-36)		28.73±2.65 (21-34)		St t=0.19	0.85
Mean ±SD (range)						
Smoking						
Yes	17	31.5	8	26.7	X ² =0.21	0.64
No	37	68.5	22	73.3		

3.2 Biochemical Assessments of endoglin

Mean endoglin in control group was 3.06±1.28 ng/ml, while it was 4.11±1.35

ng/mL in psoriasis cases. Case group showed significantly higher endoglin when compared to control group (p= 0.001) Table (2).

Table (2) Endoglin level of the studied groups.

	Case group (54)		Control group (30)		Statistical test	P value
	No	%	No	%		
Endoglin(ng/ml)	4.11±1.35 (0.48-6.6)		3.06±1.28 (0.19-6.34)		St t=3.48	0.001**
Mean ±SD (range)						

3.3 correlation between PASI score and other findings

Results of the study showed that there is positive correlation between PASI score and endoglin,

LDL,Cholesterol and fasting blood sugar with a significant value as P value (0.018,0.011,0.014 and 0.044) respectively (Table 3).

Table (3) correlation between PASI score and other findings.

PASI Score	Case group (54)	
	Correlation coefficient	P value
BMI	-0.21	0.13
Duration	-0.16	0.24
Endoglin	0.19	0.018*
Albumin	0.16	0.25
LDL	0.35	0.011*
HDL	0.09	0.54
Cholesterol	0.33	0.014*
TG	0.09	0.53
ESR 1 st hour	-0.03	0.83
ESR 2 nd hour	-0.02	0.91
CRP	-0.25	0.06
FBS	0.28	0.044*

4. Discussion

Psoriasis is a chronic, inflammatory, immune-mediated skin condition that is portrayed toward well-demarcated, erythematous plaques with silver scales, which might make connected with possibility indications of pruritus, pain, skin tightness, bleeding, Also flaking [22]. Psoriasis might show up during any age, At two peaks done agdistis onset have been reported: those Initially the middle of 20 Also 30 A long time and the second between 50 Also 60 A long time. Guys Furthermore females would just as influenced [23].

Endoglin may be An kind i film glycoprotein spotted

for Mobile surfaces Also may be and only those Transforming development factor-β (TGF-β) unpredictable. It will be predominantly communicated done endothelial units. It need likewise An urgent part Previously, angiogenesis, therefore, settling on it a paramount protein to tumor growth, survival and metastasis of tumor phones should different areas in the figure [24].

The display consider incorporated 54 patients suffice starting with psoriasis What's more with 30 clearly sound people about matched period What's more sex as a control assembly.

In the present study, there might have been no huge contrast between psoriatic patients and control groups in regards to sex. These outcomes consented for the past finding about in turn consider [25] who news person no sex predilection around psoriasis patients.

The genetic effects showed that about 18.5% from claiming psoriatic patients bring certain family history. Psoriasis need been sub ordered as stated by period of onset. Right on time onset psoriasis (also alluded should concerning illustration sort I) need onset when the period from claiming 40 years, with top onset during 16–22 a considerable length of time for age, also comprises 70% of every one psoriatic. Late-onset psoriasis, additionally termed kind ii psoriasis, demonstrates onset toward or following period 40 years, for an crest agdistis about onset the middle of 57. Also 60 a considerable length of time. An different example for HLA Acquaintanceship need been accounted. Hence, for punctual onset psoriasis, which additionally shows an solid crew history, a solid Acquaintanceship for class I HLA alleles, what's more particularly HLA-C*06, may be watched. On contrast, kind ii psoriasis will be additional sporadic also infrequently familial. Furthermore its hereditary foundation will be unclear [26]. Effects about an additional study [27] showed expanded seriousness from claiming skin lesions. Also nail love clinched alongside initial onset psoriasis for solid family history from claiming psoriasis.

Endoglin need basic part over endothelial dysfunction, hypercholesterolemia what's more atherogenesis [28]. The previous consider [29] needed news person expanded serum endoglin level on patients for hypercholesterolemia. In the different hand, another study [30], news person higher plasma level about lipids. Previously, psoriatic patients. As as much comes about uncovered a huge association between seriousness of psoriasis and serum lipid profile including serum triglyceride (TG), cholesterol, low thickness lipoprotein (LDL) what's more precise low thickness lipoprotein (VLDL). In this way move forward those comes about of the current ponder between serum level from claiming endoglin. Furthermore psoriasis.

Those gotten outcomes exhibited that there might have been helter skelter measurable critical distinction the middle of patients also controls in regards to triglycerides (TG), fasting glucose (FBS). These effects were in understanding with the individuals from claiming [17], who discovered that fasting glucose (FBS) is essentially modified done plasma starting with patients with psoriasis. What's more for [31], who discovered that level about triglycerides (TG) altogether expanded on psoriasis plasma.

Ponder about [32] led that those pervasiveness of metabolic syndrome. Previously, patients with psoriasis ranges starting with 20% with 50%, with a hazard for hosting metabolic syndrome will be no less than twofold to psoriatic patients compared for non-psoriatic control people. The Acquaintanceship between psoriasis also metabolic syndrome need essential clinical meanings and the quality about these cooperations need been over

affirmed. Eventually Tom's perusing a few observational investigations. Consequently enhance those solid affiliation the middle of psoriasis with hyperlipidemia. Also hyperglycemia which would parts from claiming metabolic syndrome.

Outcomes of the present examine demonstrated that serum endoglin levels were essentially higher. Previously, psoriatic patients. The point when contrasted with controls. Raised serum endoglin level might have been distinguished to patients with cardiovascular infections identified with endothelial brokenness. What's more hypercholesterolemia [33]. In this way raised serum endoglin level could a chance to be utilized. Likewise biomarker about hyperlipidemia. Previously, patients for psoriasis. Raised outflow for endoglin controlled. Eventually Tom's perusing immunohistochemical technique in psoriatic plaques need recently been portrayed by [34].

Those current consider demonstrated that there might have been statistically critical sure connection found between serum endoglin levels and PASI score.

With the goal extreme cases about psoriasis could be connected with large amount about serum endoglin, consequently endoglin might be a biomarker from claiming psoriasis seriousness.

Those exhibit ponder distinguished certain relationship the middle of serum endoglin level. Furthermore hdl level. On male patients of psoriasis, furthermore should sure correspondence between serum endoglin level. Also TG level over female patients for psoriasis.

Outcomes about [31] indicated critical rise for levels of downright cholesterol, triglycerides, VLDL. What's more cholesterol/HDL proportion might have been watched. Raised. In spite of might have been watched in the event about ldl and hdl levels.

5. Conclusions

From the results of present study, it is concluded that endoglin is associated with the susceptibility of development of psoriasis and has a role in its pathogenesis and severity.

6. Recommendations

The results of our study should be interpreted in light of its limitations, as the present study included a relatively small sample size. Further studies will be needed to study endoglin in other dermatological diseases. Studies on serum level of endoglin in psoriatic patients during and after local and systemic treatment are recommended. More pharmaceutical studies for finding new endoglin drug as an anti-inflammatory drug should be done.

7. Conflict Of Interest

The authors declare that they have no conflict of interest.

References

- [1] P. M.Ridker, T.Thuren, A.Zalewski, & P.Libby , Interleukin-1 β inhibition and the prevention of recurrent cardiovascular events: rationale and design of the Canakinumab Anti-inflammatory Thrombosis Outcomes Study (CANTOS). *Am Heart J*, Vol.162(4), PP..597-605,2011.
- [2] Hassan B. S, Akhtar N, kumar Paul H, Biswas C, Hossain M. A, Wahab M. A, & Zakaria A. S. M (2017): Rate of metabolic syndrome in psoriasis. *J Pak Assoc Dermatol*, Vol.27(1), PP.. 48-53,2017.
- [3] E.Burden-Teh, R.C.Phillips, K. S.Thomas, S.Ratib, D.Grindlay, & R.Murphy, A systematic review of diagnostic criteria for psoriasis in adults and children: evidence from studies with a primary aim to develop or validate diagnostic criteria. *Br J Dermatol*, Vol.178(5), PP..1035-1043,2018.
- [4] S.K.Mahil, F.Capon, & J.N.Barker , Update on psoriasis immunopathogenesis and targeted immunotherapy. In *Seminars in immunopathology*, Springer, Berlin Heidelberg , Vol. 38(1), pp. 11-27,2016.
- [5] J. B.Golden, T.S.McCormick, &N. L.Ward, IL-17 in psoriasis: implications for therapy and cardiovascular co-morbidities. *Cytokine*, Vol.62(2), PP..195–201,2013.
- [6] M.Furue, G.Tsuji, T.Chiba, & T.Kadono , Cardiovascular and metabolic diseases comorbid with psoriasis: beyond the skin. *Intern Med*, Vol.56(13), PP..1613-1619,2017.
- [7] K.Shah, L.Mellars, A.Changolkar, & S. R.Feldman, Real-world burden of comorbidities in US patients with psoriasis. *J Am Acad Dermatol*, Vol.77(2), PP..287-292,2017.
- [8] J.Takeshita, S.Grewal, S.M.Langan,N.N.Mehta, A.Ogdie, A. S.Van Voorhees, & J. M.Gelfand, Psoriasis and comorbid diseases: epidemiology. *J Am Acad Dermatol*, Vol. 76(3), PP.. 377-390,2017.
- [9] J.M.Carrascosa,V.Rocamora, R.M.Fernandez-Torres, R.Jimenez-Puya, J.C.Moreno, Coll-N.Puigserver, &E.Fonseca, Obesity and psoriasis: inflammatory nature of obesity, relationship between psoriasis and obesity, and therapeutic implications. *Actas Dermosifiliogr (English Edition)* , Vol.105 (1), pp. 31-44,2014.
- [10]M.Kunz , J.C.Simon, & A.Saalbach , Psoriasis: Obesity and Fatty Acids. *Front Immunol*, Vol.10, pp.1807,2019.
- [11]J.M.Burkhart, S.Gambaryan,S.P.Watson,K.Jurk,U.Walter,A.Sickmann,&R.P.Zahedi, What can proteomics tell us about platelets? *Circ Res*, Vol.114(7), PP..1204-1219, 2014.
- [12]F.Nassiri, M.D.Cusimano, B.W.Scheithauer, F.Rotondo, A.Fazio, G.M.Yousef, &R.V.Lloyd , Endoglin (CD105): a review of its role in angiogenesis and tumor diagnosis, progression and therapy. *Anticancer Res*; vol.31 (6), PP..2283-2290,2011.
- [13]E.Núñez-Gómez, M.Pericacho, C.Ollauri-Ibáñez, C.Bernabéu, & J.M.López-Novoa , The role of endoglin in post-ischemic revascularization. *Angiogenesis* , Vol.20(1), PP..1-24,2017.
- [14]E.Rossi, F.Sanz-Rodríguez, N.Eleno, A.Düwell, Blanco F. J, C.Langa, & C.Bernabeu , Endothelial endoglin is involved in inflammation: role in leukocyte adhesion and transmigration. *Blood*, Vol.121(2), PP.. 403-415,2013.
- [15]A.Weiss, & L.Attisano ,The TGFbeta superfamily signaling pathway. *Wiley Interdiscip Rev Dev Biol*, Vol.2(1), PP..47-63,2013.
- [16]D. Zhang, F.Zhou, X.Zhao, B.Liu, J.Chen, & J.Yang , Endoglin is a conserved regulator of vasculogenesis in zebrafish—implications for hereditary haemorrhagic telangiectasia. *Biosci Rep*, Vol.39 (5), PP..52-63,2019.
- [17]E. Gallardo-Vara, L.Ruiz-Llorente, J.Casado-Vela, M.J.Ruiz-Rodríguez, N.López-Andrés, A. K.Pattnaik, &C.Bernabeu , Endoglin Protein Interactome Profiling Identifies TRIM21 and Galectin-3 as New Binding Partners.*Cells*, Vol.8(9), PP..1082,2019.
- [18]L.Borges, V. K.Oliveira, J.Baik, S.C.Bendall, & R.C.Perlingeiro , Serial transplantation reveals a critical role for endoglin in hematopoietic stem cell quiescence. *Blood*; 133(7), 688-696,2019.
- [19]J.McDonald, P.Bayrak-Toydemir, & R.E.Pyeritz , Hereditary hemorrhagic telangiectasia: an overview of diagnosis, management, and pathogenesis. *Genet Med*, Vol.13(7), pp.607,2011.
- [20]A.M.Blázquez-Medela, L.García-Ortiz, M.A.Gómez-Marcos,J.I.Recio-Rodríguez, A.Sánchez-Rodríguez, J.M. López-Novoa, C.Martínez-Salgado , Increased plasma soluble endoglin levels as an indicator of cardiovascular alterations in hypertensive and diabetic patients. *BMC Med*, Vol.8(1), PP.. 86,2010.
- [21]D.Beiroa, A.Romero-Picó, C.Langa, C.Bernabeu, M.López, J.M.Lopez-Novoa, & C.Díeguez , Heterozygous deficiency of endoglin decreases insulin and hepatic triglyceride levels during high fat diet. , Vol.12(3), PP..52-57,2013.
- [22]T.D.Rachakonda,C.W.Schupp, & A.W.Armstrong (2014): Psoriasis prevalence among adults in the United States. *J Am Acad Dermatol*; 70(3), 512-516,2014.
- [23]D. A.Springate, R.Parisi, E. Kontopantelis, D.Reeves, C.E.M. Griffiths, &D.M.Ashcroft , Incidence, prevalence and mortality of patients with psoriasis: a UK population-based cohort study. *Br J Dermatol*, Vol.176(3), PP.. 650-658,2017.
- [24]H.Tian, J.J.Huang, C.Golzio, X.Gao, M.Hector-Greene, N.Katsanis , & G.C.Blobe , Endoglin interacts with VEGFR2 to promote angiogenesis. *The FASEB J*; Vol.32(6), PP.2934-2949,2018.
- [25]D.S.Lim, A.Bewley, H.H.Oon , Psychosocial Profile of Patients with Psoriasis. *Ann Acad Med Singapore*; Vol.47(12), PP.516-522,2018.

- [26] R. Queiro, P. Tejón, S. Alonso, & P. Coto, Age at disease onset: a key factor for understanding psoriatic disease. *Rheumatol*, Vol. 53(7), PP. 1178-1185, 2013.
- [27] P. Stuart, F. Malick, R. P. Nair, T. Henseler, H. W. Lim, S. Jenisch, J. Voorhees, E. Christophers, & J. T. Elder, Analysis of phenotypic variation in psoriasis as a function of age at onset and family history. *Arch of Dermatol Res*, Vol. 294(5), PP. 207-213, 2002.
- [28] J. Rathouska, K. Jezkova, I. Nemeckova and P. Nachtigal, Soluble endoglin, hypercholesterolemia and endothelial dysfunction. *Atherosclerosis*, Vol. 243(2), PP. 383-388, 2015.
- [29] M. Blaha, M. Cermanova, V. Blaha, P. Jarolim, C. Andrys, M. Blazek, J. Maly, L. Smolej, J. Zajic, V. Masin, & R. Zimova, Elevated serum soluble endoglin (sCD105) decreased during extracorporeal elimination therapy for familial hypercholesterolemia. *Atherosclerosis*, Vol. 197(1), PP. 264-270, 2008.
- [30] Z. HajHeydari, Serum lipids and lipoproteins in patients with psoriasis. *Arch of Iran Med*, Vol. 17(5), PP. 343, 2014.
- [31] S. Sahu, E. Devi, A. Poddar, & S. Ray, Lipid profile: a conduit in the progression from psoriasis to cardiovascular disease. *Int J Adv Med*, Vol. 4(1), PP. 173-175, 2017.
- [32] P. Gisondi, A. C. Fostini, I. Fossà, G. Girolomoni, & G. Targher, Psoriasis and the metabolic syndrome. *Clin Dermatol*, Vol. 36(1), PP. 21-28, 2018.
- [33] J. Rathouska, K. Jezkova, I. Nemeckova, L. Zemankova, M. Varejckova, & P. Nachtigal, Endoglin is not expressed with cell adhesion molecules in aorta during atherogenesis in apoE-deficient mice. *Histol Histopathol*, Vol. 30(2), PP. 233-244, 2015.
- [34] P. C. Van de Kerkhof, H. F. Rulo, J. P. van Pelt, I. M. van Vlijmen-Willems, & E. De Jong, Expression of endoglin in the transition between psoriatic uninvolved and involved skin. *Acta Derm Venereol*; 78, 19-21, 1998.